arXiv:1510.04435v1 [math.AC] 15 Oct 2015

GOLOD PROPERTY OF POWERS OF IDEALS AND OF IDEALS WITH LINEAR RESOLUTIONS

RASOUL AHANGARI MALEKI

Abstract. Let S be a regular local ring (or a polynomial ring over a field). In this paper we provide a criterion for Golodness of an ideal of S. We apply this to find some classes of Golod ideals. It is shown that for an ideal (or homogeneous ideal) $\mathfrak{a}$, there exists an integer $\rho(\mathfrak{a})$ such that for any integer $m > \rho(\mathfrak{a})$, any ideal between $\mathfrak{a}^{2m-2\rho(\mathfrak{a})}$ and $\mathfrak{a}^m$ is Golod. In the case where S is graded polynomial ring over a field of characteristic zero or where S is of dimension 2, we establish that $\rho(\mathfrak{a}) = 1$. Among other things, we prove that if an ideal $\mathfrak{a}$ is a Koszul module, then $\mathfrak{ab}$ is Golod for any ideal $\mathfrak{b}$ containing $\mathfrak{a}$.

Introduction

Let $(R, \mathfrak{m}, k)$ be a Notherian local ring (or a standard graded k-algebra) with the (homogeneous) maximal ideal $\mathfrak{m}$ and the residue field k. The Poincaré series of a finitely generated R-module M is denoted by $P^R_M(t)$ and defined to be the formal power series $\sum_{i\geq 0} \dim_k \operatorname{Tor}^R_i(M, k)t^i$. The Poincaré series $P^R_M(t)$ is rational if $P^R_M(t) = f(t)/g(t)$ for some complex polynomials $f(t)$ and $g(t)$. Rationality of a Poincaré series provides a repetitive relation for Betti numbers which can be useful in constructing a minimal free resolution. But in general this power series is not a rational function. Anick [1] discovered the first example of a local ring R such that $P^R_k(t)$ is not a rational function. Also see [14] for more such examples. However counterexamples do not seem to be plentiful.

Let $(S, \mathfrak{n}, k)$ and $(R, \mathfrak{m}, k)$ are Noetherian local rings (or a standard graded k-algebra) with the maximal (or homogeneous maximal) ideals $\mathfrak{n}$ and $\mathfrak{m}$ respectively, and with the same residue field k. Let $\varphi : (S, \mathfrak{n}, k) \to (R, \mathfrak{m}, k)$ be a surjective ring homomorphism. Then there is a coefficientwise inequality of formal power series which was initially derived by Serre :

$$P^R_k(t) \preceq \frac{P^S_k(t)}{1 - t(P^S_R(t) - 1)}.$$

The homomorphism φ is said to be Golod if the equality holds. In the case where S is a regular local ring (or a polynomial over k) and $\dim S = \operatorname{embdim} R$ and $\mathfrak{a} = \ker \varphi$ we say that R is Golod, or the ideal $\mathfrak{a}$ is Golod, if the homomorphism φ is Golod . In this case the Golodness of R implies that $P^R_k(t)$ is rational. More than this, Golod rings are an example

2010 *Mathematics Subject Classification.* 13A02, 13D02, 13H02 .

Key words and phrases. Powers of ideals, Golod rings; linear resolutions.

This research was in part supported by a grant from IPM (No. 94130028).

of good rings, in the sense that all finitely generated modules over such rings have rational Poincaré series sharing a common denominator [2]. In the case where S is a polynomial ring over a field of characteristic zero, Herzog and Huneke [7] find quite large classes of Golod ideals. They show that for a homogeneous ideal $\mathfrak{a}$, the ideals $\mathfrak{a}^m$, $\mathfrak{a}^{(m)}$ (the m-th symbolic power of $\mathfrak{a}$) and $\widetilde{\mathfrak{a}^m}$ (the saturated power of $\mathfrak{a}$) are Golod for all $m \geq 2$. Their proofs hinge on the definition of strongly Golod ideals. The authors call an ideal $\mathfrak{a}$ is strongly Golod if $\partial(\mathfrak{a})^2 \subseteq \mathfrak{a}$. Here $\partial(\mathfrak{a})$ denotes the ideal generated by all the partial derivatives of elements of $\mathfrak{a}$. They show that strongly Golod ideals are Golod.

In view of these results, it is a natural expectation that the same results of [7] also must be true when S is a regular local ring (or a polynomial ring over a field of any characteristic). A known fact in this direction is a result of Herzog, Welker and Yassemi [10] which states that large powers of an ideal are Golod. Also in [8] it is shown that if $\mathfrak{a}$ and $\mathfrak{b}$ are ideals of a regular local ring (or a polynomial ring over a field) and $\mathfrak{a} \cap \mathfrak{b} = \mathfrak{a}\mathfrak{b}$, then $\mathfrak{a}\mathfrak{b}$ is Golod. Viewing these known Golod ideals, another ideals which may candidate for being Golod are products of ideals. Newly Stefani [20] find an example of two monomial ideals in a polynomial ring over a field, whose product is not Golod.

In this paper we are going to find some new classes of Golod ideals of a regular local ring ((or polynomial ring over a filed). In Section one we show the following: A surjective homomorphism $\varphi : (S, \mathfrak{n}, k) \to (R, \mathfrak{m}, k)$ of local rings is Golod if there exists a proper ideal L of R satisfying $L^2 = 0$ and the induced maps

$$\operatorname{Tor}_i^S(R, k) \to \operatorname{Tor}_i^S(R/L, k)$$

by the projection $R \to R/L$ are zero for all $i > 0$. In the case where S is a regular local ring (or a polynomial ring over a field), this provides a criterion for Golodness of an ideal of S. In section two we apply this to get some class of Golod ideals.

Let $\mathfrak{a}$ and $\mathfrak{b}$ be proper ideals of a regular local ring (or a polynomial ring over k) $(S, \mathfrak{n}, k)$. We show that there exists a positive integer $\rho(\mathfrak{a})$ (see Section 2 for the definition) such that any ideal between $\mathfrak{a}^{2(m-\rho(\mathfrak{a}))}$ and $\mathfrak{a}^m$ is Golod for all $m > \rho(\mathfrak{a})$. In the following cases we are able to prove that $\rho(\mathfrak{a}) = 1$.

(1) S is a polynomial ring over a field of characteristic zero, and $\mathfrak{a}$ any homogeneous ideal of S;
(2) S has Krull dimension at most 2;
(3) $\mathfrak{a}$ is generated by a part of a regular system of parameter of S.

Also we show that if $\mathfrak{a}$ is a Koszul ideal (that is the ideal whose associated graded module with respect to $\mathfrak{n}$ has a linear resolution) then $\mathfrak{a}\mathfrak{b}$ is Golod for all ideal $\mathfrak{b}$ containing $\mathfrak{a}$, see Theorem 2.10. In particular case if $\mathfrak{a}$ generated by a regular system of parameter and $\mathfrak{b}$ contains a power $\mathfrak{a}^r$, then $\mathfrak{a}^r\mathfrak{b}$ is Golod.

1. Golod homomorphisms and Massey operations

There is an important tool for investigating Golodness of a surjective homomorphism of local rings and studying of resolutions. We use this tool in this section.

Let $\varphi : (S, \mathfrak{n}, k) \to (R, \mathfrak{m}, k)$ be a surjective homomorphism of local rings. Assume $\mathcal{D}$ is a minimal free resolution of k over S equipped with a graded commutative DG-algebra structure; such a resolution always exists, see [16]. Let $\mathcal{A} = \mathcal{D} \otimes_S R$. Then $\mathcal{A}$ is a graded commutative DG-algebra. We denote $Z(\mathcal{A})$, $B(\mathcal{A})$ and $\mathrm{H}(\mathcal{A}) = \frac{Z(\mathcal{A})}{B(\mathcal{A})}$ the module of cycles, boundaries and homologies of $\mathcal{A}$ respectively. If a is a homogeneous element of $\mathcal{A}$, the degree of a is denoted by $|a|$ and we set $\bar{a} = (-1)^{|a|+1}a$.

According to Gulliksen, we say $\mathcal{A}$ admits a trivial Massey operation if for some homogeneous k-basis $\mathcal{B} = \{h_i\}_{\geq 1}$ of $\mathrm{H}_{\geq 1}(\mathcal{A}) := \bigoplus_{i\geq 1} \mathrm{H}_i(\mathcal{A})$ there exists a function $\mu : \bigsqcup_{n=1}^{\infty} \mathcal{B}^n \to \mathcal{A}$, such that

(1) $\mu(h_\lambda) = z_\lambda \in Z(\mathcal{A})$ with $cls(z) = h$;

(2) $\partial\mu(h_{\lambda_1}, \cdots, h_{\lambda_n}) = \sum_{j=1}^{n-1} \overline{\mu(h_{\lambda_1}, \cdots, h_{\lambda_j})}\mu(h_{\lambda_{j+1}}, \cdots, h_{\lambda_n}) \quad \text{for } n \geq 2$;

(3) $\mu(\mathcal{B}^n) \subseteq \mathfrak{m}\mathcal{A} \quad \text{for } n \geq 1$.

It is well known that the homomorphism φ is Golod if and only if the *DG* algebra $\mathcal{A}$ admits a trivial Massey operation (see [2] and [5]).

The following provides a criterion for Golodness of a homomorphism. The idea of this was motivated by [15, Lemma 1.2]. We apply similar technic for the proof.

Lemma 1.1. *If there exists a proper ideal L of R with $L^2 = 0$ such that the map*

$$\mathrm{Tor}_i^S(R, k) \to \mathrm{Tor}_i^S(R/L, k)$$

induced by the projection $R \to R/L$ is zero for all $i > 0$, then the map φ is Golod. Moreover, the Massey operation μ can be constructed so that $\operatorname{Im}\mu \subseteq L\mathcal{A}$.

Proof. Let $\mathcal{D}$ and $\mathcal{A} = \mathcal{D} \otimes_S R$ be as above. For proving that φ is Golod we show that μ can be chosen that $\mu(h_{\lambda_1})\mu(h_{\lambda_2}) = 0$ for all $h_{\lambda_1}, h_{\lambda_2} \in \mathcal{B}$.

We have the isomorphisms $\mathcal{D} \otimes_S R/L \cong (\mathcal{D} \otimes_S R) \otimes_R R/L = \mathcal{A} \otimes_R R/L \cong \mathcal{A}/L\mathcal{A}$ of complexes of S-modules. Hence the map $\mathrm{Tor}_i^S(R, k) \to \mathrm{Tor}_i^S(R/L, k)$ can be identified with the map

$$\psi_i : \mathrm{H}_i(\mathcal{A}) \to \mathrm{H}_i(\mathcal{A}/L\mathcal{A})$$

induced by the projection $\mathcal{A} \to \mathcal{A}/L\mathcal{A}$. Now let $h_\lambda \in \mathcal{B}$ so $h_\lambda = cls(z)$ for some $z \in Z_i(\mathcal{A})$ and for some $i > 0$. Since $\psi_i(h_{z_\lambda}) = 0$, there is an element $x \in B_i(\mathcal{A})$ such that $z - x \in L\mathcal{A}_i$. By setting $z_\lambda = z - x$ which is a cycle, we have $h_\lambda = cls(z_\lambda)$. Therefore every element h_λ of $\mathcal{B}$ can be represented as $cls(z_\lambda)$ for some $z_\lambda \in Z(\mathcal{A}) \cap L\mathcal{A}$. Now we define $\mu(h_\lambda) = z_\lambda$. Since $L^2 = 0$, for any two element $h_{\lambda_1}, h_{\lambda_2}$ we get $\mu(h_{\lambda_1})\mu(h_{\lambda_2}) = 0$. By using this property

we may set $\mu(h_{\lambda_1}, \cdots, h_{\lambda_n}) = 0$ for all $n \geq 2$ and then obviously (2) is satisfied and φ is Golod. □

In the rest of the paper $(S, \mathfrak{n}, k)$ denotes a regular local ring (or a polynomial over k) with the maximal (or homogeneous maximal) ideal $\mathfrak{n}$ and the residue field k. In the graded case all modules considered to be graded. Also we set $d = \dim S$.

Remark 1.2. *Assume that K is the Koszul complex of S with respect to a minimal system of generators of $\mathfrak{n}$. We denote by $\mathcal{Z}$ the cycles of K. The complex K is a minimal free resolution of k. For ideals $\mathfrak{a} \subseteq \mathfrak{b}$ of S we have the commutative diagram*

$$\begin{array}{ccc} \operatorname{Tor}_i^S(S/\mathfrak{a}, k) & \xrightarrow{\varphi_i} & \operatorname{Tor}_i^S(S/\mathfrak{b}, k) \\ \downarrow \cong & & \downarrow \cong \\ \operatorname{Tor}_{i-1}^S(\mathfrak{a}, k) & \longrightarrow & \operatorname{Tor}_{i-1}^S(\mathfrak{b}, k) \\ \downarrow \cong & & \downarrow \cong \\ \mathrm{H}_{i-1}(\mathfrak{a}K) & \longrightarrow & \mathrm{H}_{i-1}(\mathfrak{b}K) \end{array} \tag{1}$$

for all $i \geq 1$, where the top row induced by the natural isomorphism $S/\mathfrak{a} \to S/\mathfrak{b}$, the middle row by the inclusion $\mathfrak{a} \subseteq \mathfrak{b}$ and the bottom row by the inclusion $\mathfrak{a}K \subseteq \mathfrak{b}K$ (of complexes).

The following provides a criterion for the Golodness of an ideal.

Proposition 1.3. *Let the situation be as Remark 1.2. Assume that $\mathfrak{b}^2 \subseteq \mathfrak{a} \subseteq \mathfrak{b}$ and one of the following equivalent conditions hold.*

(1) $\mathcal{Z}_i \cap \mathfrak{a}K_i \subseteq \mathfrak{b}\mathcal{Z}_i$ for all $i \geq 1$;
(2) $\operatorname{Tor}_i^S(S/\mathfrak{a}, k) \to \operatorname{Tor}_i^S(S/\mathfrak{b}, k)$ is zero for all $i \geq 1$.

Then the ideal $\mathfrak{a}$ is a Golod.

Proof. For an ideal $\mathfrak{c}$ of S we have

$$\mathrm{H}_i(\mathfrak{c}K) = \mathrm{H}(\mathfrak{c}K_{i+1} \to \mathfrak{c}K_i \to \mathfrak{c}K_{i-1}) = \frac{\mathcal{Z}_i \cap \mathfrak{c}K_i}{\mathfrak{c}\mathcal{Z}_i}.$$

Hence in view of Remark 1.2, for each $i \geq 1$, the map $\operatorname{Tor}_i^S(S/\mathfrak{a}, k) \to \operatorname{Tor}_i^S(S/\mathfrak{b}, k)$ can be identified with the natural map

$$\frac{\mathcal{Z}_i \cap \mathfrak{a}K_i}{\mathfrak{a}\mathcal{Z}_i} \to \frac{\mathcal{Z}_i \cap \mathfrak{b}K_i}{\mathfrak{b}\mathcal{Z}_i}.$$

Thus (1) and (2) are equivalent. Set $R := S/\mathfrak{a}$ and $L := \mathfrak{b}/\mathfrak{a}$. For every $i \geq 1$, the map

$$\operatorname{Tor}_i^S(R, k) \to \operatorname{Tor}_i^S(R/L, k)$$

can be identified with the map

$$\operatorname{Tor}_i^S(S/\mathfrak{a}, k) \to \operatorname{Tor}_i^S(S/\mathfrak{b}, k)$$

which is zero. Also by the hypothesis we get $L^2 = 0$. Now the assertion follows from Lemma 1.1. □

2. Golod ideals

In this section we apply the results of previous section to obtain some classes of Golod ideals.

Assume that $\mathfrak{c}$ is an ideal of S and N is a submodule of finitely generated S-module M. Artin-Rees lemma [4, Lemma 5.1] states that there exists an integer r such that

$$N \cap \mathfrak{c}^m M = \mathfrak{c}^{m-r}(N \cap \mathfrak{c}^r M)$$

for all $m \geq r$. The smallest such number r is called the Artin-Rees number. A slightly weaker statement which follows from the lemma, and that is good enough for application, is that there exists a positive integer r such that for all $m \geq r$ the following inclusion holds

$$N \cap \mathfrak{c}^m M \subseteq \mathfrak{c}^{m-r} N.$$

Following the notation of Section one let K and $\mathcal{Z}$ be the Koszul complex and Koszul cycles. Thus from the above argument, for an ideal $\mathfrak{c}$ of S there exists the smallest integer $\rho_i(\mathfrak{c})$ such that the inclusion

$$\mathcal{Z}_i \cap \mathfrak{c}^m K_i \subseteq \mathfrak{c}^{m-\rho_i(\mathfrak{c})} \mathcal{Z}_i$$

holds for all $m \geq \rho_i(\mathfrak{c})$. Define $\rho(\mathfrak{c})$ to be the number $\max\{\rho_1(\mathfrak{c}), \cdots, \rho_d(\mathfrak{c})\}$. We call this number the *Koszul Artin-Ress* number of the ideal $\mathfrak{c}$.

Remark 2.1. *It follows from Remark 1.2 that $\rho(\mathfrak{c})$ is the smallest integer n such that for all $m \geq n$ and $i \geq 1$, the maps*

$$\operatorname{Tor}_i^S(S/\mathfrak{c}^m, k) \to \operatorname{Tor}_i^S(S/\mathfrak{c}^{m-n}, k)$$

are zero.

The following theorem covers a result of Herzog et al. [10] which says that all higher powers of an ideal of a regular local ring are Golod.

Theorem 2.2. *Let $\mathfrak{c}$ be an ideal of S and m be a positive integer with $m > \rho(\mathfrak{c})$. If $\mathfrak{a}$ is an ideal of S such that $\mathfrak{c}^{2(m-\rho(\mathfrak{c}))} \subseteq \mathfrak{a} \subseteq \mathfrak{c}^m$. Then $\mathfrak{a}^p$ is Golod for all $p \geq 1$. In particular $\mathfrak{c}^m$ is Golod.*

Proof. Let $p \geq 1$ be an integer. We have the following inclusions

$$\mathcal{Z}_i \cap \mathfrak{a}^p K_i \subseteq \mathcal{Z}_i \cap \mathfrak{c}^{mp} K_i \subseteq \mathfrak{c}^{mp-\rho(\mathfrak{c})} \mathcal{Z}_i$$

for all $1 \leq i \leq d$ where the right inclusion follows from the definition of $\rho(\mathfrak{c})$. Set $\mathfrak{b} := \mathfrak{c}^{mp-\rho(\mathfrak{c})}$ thus we get $\mathfrak{b}^2 \subseteq \mathfrak{a}^p \subseteq \mathfrak{b}$ and $\mathcal{Z}_i \cap \mathfrak{a}^p K_i \subseteq \mathfrak{b}\mathcal{Z}_i$. Now applying Proposition 1.3 we conclude that $\mathfrak{a}^p$ is a Golod ideal. □

Corollary 2.3. *Let $\mathfrak{a}$, $\mathfrak{b}$ and $\mathfrak{c}$ be ideals of S. Assume that for some integers $p, q \geq \rho(\mathfrak{c})$ the following containments hold*

$$\mathfrak{c}^{2p-\rho(\mathfrak{c})} \subseteq \mathfrak{a} \subseteq \mathfrak{c}^p, \quad \mathfrak{c}^{2q-\rho(\mathfrak{c})} \subseteq \mathfrak{b} \subseteq \mathfrak{c}^q.$$

Then $\mathfrak{ab}$ is Golod.

Proof. From the hypothesis we have $\mathfrak{c}^{2(p+q-\rho(\mathfrak{c}))} \subseteq \mathfrak{ab} \subseteq \mathfrak{c}^{p+q}$. Now by the above theorem we get the desired. □

For an ideal $\mathfrak{c}$ of S we have $\rho(\mathfrak{c}) \geq 1$, by definition. In view of Theorem 2.2, it would be good if $\rho(\mathfrak{c}) = 1$. In the graded case we show that the Koszul Artin-Ress number of any homogeneous ideal reaches its lower bound.

Remark 2.4. *Let $S = k[X_1, \cdots, X_d]$ be a graded polynomial ring over a field k of characteristic zero. For a homogeneous ideal $\mathfrak{c}$ of S with $\mathfrak{c} \subseteq (X_1, \cdots, X_d)^2$, let*

$$0 \to F_n \xrightarrow{\phi_n} F_{n-1} \to \cdots \to F_1 \xrightarrow{\phi_1} F_0 \to S/\mathfrak{c} \to 0$$

be the graded minimal free resolution of $S/\mathfrak{c}$. Let b_i be the rank of F_i and $f_{i1}, \cdots, f_{ib_i}$ be a homogeneous basis of F_i. Also, assume that

$$\phi_i(f_{ij}) = \sum_{k=1}^{b_{i-1}} \alpha_{jk}^{(i)} f_{i-1k},$$

where the $\alpha_{jk}^{(i)}$ are homogeneous polynomials in S. Set $R = S/\mathfrak{c}$ and let K^R be the Koszul complex of the ring R with respect to a minimal homogeneous generating set of the graded maximal ideal of the ring. Then K_1^R is the free module $\oplus_{i=1}^d Re_i$ with the basis $e_1, \cdots, e_d$ and for each $l = 1, \cdots, d$ the elements $e_{i_1} \wedge \cdots \wedge e_{i_l}$ provide the natural R-basis for the free module $K_l^R = \bigwedge^l (\oplus_{i=1}^d Re_i)$. From a result of Herzog [6, Corollary 2] *for each $l = 1, \cdots, n$ a homogeneous k-basis of $\mathrm{H}_l(K^R)$ is given by cycles of the form*

$$z = \sum_{1 \leq i_1 < i_2 < \cdots < i_l \leq d} u_{j_1, \cdots, j_l} e_{i_1} \wedge \cdots \wedge e_{i_l}$$

where each $u_{j_1, \cdots, j_l}$ is a linear combinations of Jacobians of the form

$$\frac{\partial(\alpha_{j_1 j_2}^{(l)}, \alpha_{j_2 j_3}^{(l-1)}, \cdots, \alpha_{j_l 1}^{(1)})}{\partial(x_{i_1}, \cdots, x_{i_l})}$$

with $1 \leq j_k \leq b_{l-k+1}$. Here the Jacobian $\frac{\partial(g_1, \cdots, g_l)}{\partial(x_{i_1}, \cdots, x_{i_l})}$ of the polynomials $g_1, \cdots, g_l$ with respect to $x_{i_1}, \cdots, x_{i_l}$ is defined to be

$$\det\left(\frac{\partial g_k}{\partial X_{i_j}}\right)_{1 \leq k, j \leq l} \ mod \ \ \mathfrak{c}.$$

Denote by $\partial(\mathfrak{c})$ the ideal generated by partial derivatives $\partial f / \partial X_i$ with $f \in \mathfrak{c}$ and $i = 1, \cdots, d$. Since the elements $\alpha_{j_l 1}^{(1)}$ with $j_l = 1, \cdots, b_1$ generate $\mathfrak{c}$, we see that $u_{j_1, \cdots, j_l} \in \partial(\mathfrak{c})$. Thus the

homology classes of elements of $\mathcal{Z}_l^R \cap \partial(\mathfrak{c})K_l^R$ *generate the homology module* $\mathrm{H}_l(K^R)$*, where* $\mathcal{Z}_l^R$ *denotes the* l*-cycles of* K^R*.*

Theorem 2.5. *Let* S *be a graded polynomial ring over a field of characteristic zero. Then for any homogeneous proper ideal* $\mathfrak{a}$ *of* S *the following statement hold.*

(*i*) $\rho(\mathfrak{a}) = 1$*;*

(*ii*) *any homogeneous ideal* $\mathfrak{b}$ *with* $\mathfrak{a}^{2m-2} \subseteq \mathfrak{b} \subseteq \mathfrak{a}^m$ *is Golod.*

Proof. We first prove the following claim: if $\mathfrak{c}_1 \subseteq \mathfrak{c}_2$ are homogeneous ideals of S such that $\partial(\mathfrak{c}_1) \subseteq \mathfrak{c}_2$, then the map $\mathrm{Tor}_i^S(S/\mathfrak{c}_1, k) \to \mathrm{Tor}_i^S(S/\mathfrak{c}_2, k)$ is zero for all $i > 0$. Set $R_1 = S/\mathfrak{c}_1$ and $R_2 = S/\mathfrak{c}_2$. Let $e_1, \cdots, e_d$ and $f_1, \cdots, f_d$ be the natural basis of the free modules $K_1^{R_1} = \oplus_{i=1}^d R_1 e_i$ and $K_1^{R_2} = \oplus_{i=1}^d R_2 f_i$ respectively. Consider the natural morphism of complexes

$$\psi : K^{R_1} \to K^{R_2}$$

where $\psi((\mathfrak{c}_1 + 1)e_i) = (\mathfrak{c}_2 + 1)f_i$ for all $i = 1, \cdots, d$. For any homogeneous ideal $\mathfrak{c}$ of S there is a natural isomorphism $S/\mathfrak{c} \otimes_S K^S \to K^{S/\mathfrak{c}}$ of complexes. Thus the map $\mathrm{Tor}_i^S(R_1, k) \to \mathrm{Tor}_i^S(R_2, k)$ can be identified with the natural map

$$\mathrm{H}_i(\psi) : \mathrm{H}_i(K^{R_1}) \to \mathrm{H}_i(K^{R_1})$$

induced by ψ on homology modules. From the above remark, the homology classes of elements of $\mathcal{Z}_i^{R_1} \cap \partial(\mathfrak{c}_1)K_i^{R_1}$ generate $\mathrm{H}_i(K^{R_1})$. Now since $\partial(\mathfrak{c}_1) \subseteq \mathfrak{c}_2$ we have $\psi_i(\mathcal{Z}_i^{R_1} \cap \partial(\mathfrak{c}_1)K_i^{R_1}) \subseteq \partial(\mathfrak{c}_1)K_i^{R_2} = 0$. Thus $\mathrm{H}_i(\psi) = 0$ and this complete the proof of the claim. For (i), applying we Remark 2.1 and Theorem 2.2, it is enough to show that the map

$$\mathrm{Tor}_i^S(S/\mathfrak{a}^m, k) \to \mathrm{Tor}_i^S(S/\mathfrak{a}^{m-1}, k)$$

is zero for all $i > 0$. To this end, observe that $\partial(\mathfrak{a}^m) \subseteq \partial(\mathfrak{a})\mathfrak{a}^{m-1}$ and apply the claim with $\mathfrak{c}_1 = \mathfrak{a}^m$ and $\mathfrak{c}_2 = \mathfrak{a}^{m-1}$. Using Theorem 2.2, (ii) is a direct consequence of (i). □

Motivated by the above theorem we ask the following natural question.

Question 2.6. *Let* S *be a regular local ring. Is it true that* $\rho(\mathfrak{a}) = 1$ *for any proper ideal* $\mathfrak{a}$ *of* S *or equivalently that the map*

$$\mathrm{Tor}_i^S(S/\mathfrak{a}^m, k) \to \mathrm{Tor}_i^S(S/\mathfrak{a}^{m-1}, k)$$

is zero for all $i > 0$*?*

At least in the case where $\dim S \leq 2$ the answer is positive. The case that $\dim S = 1$ is obvious. The following is for the case of dimension two.

Theorem 2.7. *Let* $\mathfrak{a}$ *be an ideal of the regular local ring (or a polynomial ring over a field)* S *of dimension* 2*. Then* $\rho(\mathfrak{a}) = 1$*.*

Proof. Let $m > 0$. We show that the map

$$\mathrm{Tor}_i^S(S/\mathfrak{a}^m, k) \to \mathrm{Tor}_i^S(S/\mathfrak{a}^{m-1}, k)$$

is zero for all $i > 0$. For the case where $i = 1$ the map is identified with the natural map $\mathfrak{n} \cap \mathfrak{a}^m/\mathfrak{n}\mathfrak{a}^m \to \mathfrak{n} \cap \mathfrak{a}^{m-1}/\mathfrak{n}\mathfrak{a}^{m-1}$ which is clearly zero.

For a sequence $x_1, \cdots, x_t$ of elements of S and for all S-module M, we have the following exact sequences

$$0 \to \mathrm{H}_0(K(x_t) \otimes \mathrm{H}_i(x_1, \cdots, x_{t-1}; M)) \to \mathrm{H}_i(x_1, \cdots, x_t; M)$$
$$\to \mathrm{H}_1(K(x_t) \otimes \mathrm{H}_{i-1}(x_1, \cdots, x_{t-1}; M)) \to 0$$

of Koszul homology modules, see [19]. Now let x, y be a regular system of parameter of S then by replacing M with $S/\mathfrak{a}^m$ and $S/\mathfrak{a}^{m-1}$ we get the commutative diagram
(2)

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \mathrm{H}_0(K(y) \otimes \mathrm{H}_2(x; S/\mathfrak{a}^m)) & \longrightarrow & \mathrm{H}_2(x, y; S/\mathfrak{a}^m) & \longrightarrow & \mathrm{H}_1(K(y) \otimes \mathrm{H}_1(x; S/\mathfrak{a}^m)) & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle \alpha} & & \downarrow{\scriptstyle \beta} & & \downarrow{\scriptstyle \gamma} & & \\
0 & \longrightarrow & \mathrm{H}_0(K(y) \otimes \mathrm{H}_2(x; S/\mathfrak{a}^{m-1})) & \longrightarrow & \mathrm{H}_2(x, y; S/\mathfrak{a}^{m-1}) & \longrightarrow & \mathrm{H}_1(K(y) \otimes \mathrm{H}_1(x; S/\mathfrak{a}^{m-1})) & \longrightarrow & 0
\end{array}$$

We have to show that $\beta = 0$. Since $\mathrm{H}_2(x; S/\mathfrak{a}^m) = 0 = \mathrm{H}_2(x; S/\mathfrak{a}^{m-1})$, we need to show that $\gamma = 0$. The map γ is induced by the map $\mathrm{H}_1(x; S/\mathfrak{a}^m) \to \mathrm{H}_1(x; S/\mathfrak{a}^{m-1})$ where can be identified with the map

$$xS \cap \mathfrak{a}^m/x\mathfrak{a}^m \to xS \cap \mathfrak{a}^{m-1}/x\mathfrak{a}^{m-1}.$$

Therefore it is enough to show that $(\mathfrak{a}^m : x) \subseteq \mathfrak{a}^{m-1}$. Note that the ring S/xS is a regular local of dimension one and so the image of $\mathfrak{a}$ in S/xS is generated by an element $xS + u$ for some $u \in \mathfrak{a}$. Now it is easy to see that there are elements $r_1, \cdots, r_t$ in S such that $\mathfrak{a} = (u, r_1x, \cdots, r_tx)$. One has $\mathfrak{a}^m \subseteq Su^m + x\mathfrak{a}^{m-1}$. If $z \in (\mathfrak{a}^m : x)$, then we can write $zx = su^m + bx$ for some $s \in S$ and $b \in \mathfrak{a}^{m-1}$ and we have $(z - b)x = su^m$. Since (x) is a prime ideal of S we get $u \in (x)$ or $s \in (x)$. In any case we can obtain that $z \in \mathfrak{a}^{m-1}$. Therefore $(\mathfrak{a}^m : x) \subseteq \mathfrak{a}^{m-1}$. □

2.1. **Golodness of ideals with linear resolutions.** Let A be a standard graded algebra over a field k and N be a graded A-module with a minimal generating set all of the same degree q. We say that N has a q-linear resolution if $\mathrm{Tor}_i^A(N, k)_j = 0$ for all i and all $j \neq i+q$. Also, we say that N is componentwise linear if for all integer q the graded submodule $N_{\langle q \rangle}$ generated by all homogeneous elements of N with degree q, has a q-linear resolution.

There is an analogue of the notion of modules with linear resolution which is defined in both local and graded case. Let $(R, \mathfrak{m}, k)$ be a local ring (or a standard graded k-algebra) with the maximal (or homogeneous maximal)ideal $\mathfrak{m}$. An R-module M is called Koszul if its associated graded module $\mathrm{gr}_{\mathfrak{m}}(M) = \oplus_{i \geq 0}\mathfrak{m}^iM/\mathfrak{m}^{i+1}M$ as a graded $gr_{\mathfrak{m}}(R)$-module has linear resolution. If the residue field k is Koszul we say that the ring R is Koszul. Note that in the graded case $gr_{\mathfrak{m}}(R)$ is identified with R itself and any graded module with linear resolution

is a Koszul module. However such a graded modules are not the only modules which are Koszul see [9, example 1.9]. If R is a graded Koszul algebra there is a characterization of (graded) Koszul modules due to Römer [17]: A graded R-module M is Koszul if and only if M is componentwise linear.

Let S be a polynomial ring with standard grading. It is known [3] that every graded ideals of S with linear resolution is Golod. This result generalized by Herzog, Reiner and Welker for componentwise linear ideals of S. Since S is a Koszul algebra, in view of the characterization of Römer this can be restated in the following form: any Koszul ideal of S is Golod. Motivated by this the following natural question raised in the local case.

Question 2.8. *Let S be a regular local ring. Is any Koszul ideal of S Golod?*

Unfortunately we do not have an answer to this question, but in what follows we show some relations between Golod ideals and Koszul ideals.
There is a characterization, due to Şega, of Koszul module. This provides a necessary condition for Koszulness of a module.

Remark 2.9. *(see* [18, Theorem 2.2 (c)]*) Let $(R, \mathfrak{m}; k)$ be a local ring (or a standard graded k-algebra). If an R-module M is Koszul, then the map*

$$\operatorname{Tor}_i^R(M, R/\mathfrak{m}^2) \to \operatorname{Tor}_i^R(M, R/\mathfrak{m})$$

is zero for all $i > 0$.

In the graded case, when M generated by elements of the same degree, one can see that this condition is equivalent to say that M has a linear resolution.

Theorem 2.10. *Let $\mathfrak{a}$ and $\mathfrak{b}$ be ideals of S such that $\mathfrak{a} \subseteq \mathfrak{b}$. If the map*

$$\delta_i : \operatorname{Tor}_i^S(\mathfrak{a}, S/\mathfrak{n}^2) \to \operatorname{Tor}_i^S(\mathfrak{a}, S/\mathfrak{n})$$

induced by the natural projection $S/\mathfrak{n}^2 \to S/\mathfrak{n}$ is zero for all $i > 0$, then $\mathfrak{ab}$ is Golod. In particular, if $\mathfrak{a}$ is Koszul (as an S-module), then $\mathfrak{ab}$ is Golod.

Proof. Applying Proposition 1.3, and Remark 1.2 it is enough to show that the map

$$\alpha_i : \operatorname{Tor}_i^S(\mathfrak{ab}, S/\mathfrak{n}) \to \operatorname{Tor}_i^S(\mathfrak{a}, S/\mathfrak{n})$$

is zero for all $i > 0$. Using the exact sequence $0 \to \mathfrak{n}/\mathfrak{n}^2 \to S/\mathfrak{n}^2 \to S/\mathfrak{n} \to 0$ we get the following commutative diagram

$$\begin{array}{ccccc} & & \operatorname{Tor}_i^S(\mathfrak{ab}, S/\mathfrak{n}) & \longrightarrow & \operatorname{Tor}_{i-1}^S(\mathfrak{ab}, \mathfrak{n}/\mathfrak{n}^2) \\ & & \downarrow{\scriptstyle \alpha_i} & & \downarrow{\scriptstyle \beta_{i-1}} \\ \operatorname{Tor}_i^S(\mathfrak{a}, S/\mathfrak{n}^2) & \xrightarrow{\delta_i} & \operatorname{Tor}_i^S(\mathfrak{a}, S/\mathfrak{n}) & \xrightarrow{\gamma_i} & \operatorname{Tor}_{i-1}^S(\mathfrak{a}, \mathfrak{n}/\mathfrak{n}^2) \end{array} \tag{3}$$

for any $i > 0$. For all $i > 0$, the map γ_i is injective since $\delta_i = 0$ by the hypothesis. This in conjunction with the fact that $\mathfrak{n}/\mathfrak{n}^2$ is a $S/\mathfrak{n}$-vector space implies that $\alpha_i = 0$ if $\beta_{i-1} = 0$.

One can see that $\beta_0 = 0$ and so $\alpha_1 = 0$. Observe that β_i is a direct sum of α_i. Therefore using induction on i, one concludes the desired. In the particular case where $\mathfrak{a}$ is Koszul by the above remark $\delta_i = 0$ for all $i > 0$ thus we get the conclusion. □

We remark that in the above theorem the condition $\mathfrak{a} \subseteq \mathfrak{b}$ is necessary: Let k be a field and $S = k[X, Y, Z, W]$ a polynomial ring. Then homogeneous maximal ideal $\mathfrak{a} = (X, Y, Z, W)$ is Koszul. Consider the ideal $\mathfrak{b} = (X^2, Y^2, Z^2, W^2)$. The the ideal $\mathfrak{ab}$ is not Golod, see [20, Example 2.1].

It is known that the powers of maximal ideal of a Koszul local ring are Koszul modules. When the ring is regular, this result can be extend to the powers of an ideal generated by a part of a regular system of parameter.

Lemma 2.11. *Let $\mathfrak{p}$ be an ideal of S generated by a part of a regular system of parameter. Then $\mathfrak{p}^r$ is a Koszul module.*

Proof. Let $x_1, \cdots, x_u$ be a part of a regular system of parameter and $\mathfrak{p} = (x_1, \cdots, x_u)$. Extend this sequence to a regular system of parameter $x_1, \cdots, x_u, x_{u+1}, \cdots x_v$, where $v = \dim_k \mathfrak{n}/\mathfrak{n}^2$. Set $\mathfrak{q} := (x_{u+1}, \cdots x_v)$, $\bar{S} := S/\mathfrak{q}$ and $\bar{\mathfrak{n}} := \mathfrak{n}/\mathfrak{q}$. $\bar{S}$ is a regular local ring an then the module $\mathfrak{p}^r + \mathfrak{q}/\mathfrak{q} \cong \bar{\mathfrak{n}}^r$ is Koszul over $\bar{S}$. Set $x^* = x + \mathfrak{n}^2$ for any x in $\mathfrak{n} \setminus \mathfrak{n}^2$. Since $x^*_{u+1}, \cdots, x^*_v$ is a regular sequence of degree one on $\mathrm{gr}_{\mathfrak{n}}(S)$ and $\mathrm{gr}_{\mathfrak{n}}(\bar{S}) = \mathrm{gr}_{\mathfrak{n}}(S)/(x^*_{u+1}, \cdots, x^*_v)$, we see that $\mathrm{gr}_{\mathfrak{n}}(\bar{S})$ has a linear resolution as a graded $\mathrm{gr}_{\mathfrak{n}}(S)$-module. Therefore $\bar{S}$ is a Koszul S-module. Now by [12, Theeorem 5.2], $\mathfrak{p}^r + \mathfrak{q}/\mathfrak{q}$ is a Koszul S-module. Also one can see that $x^*_{u+1}, \cdots, x^*_v$ forms a regular sequence on $\mathrm{gr}_{\mathfrak{n}}(S)$-module $\mathrm{gr}_n(S/\mathfrak{p}^r)$. Applying [11, Theorem 2.13 (c)], we conclude that $\mathfrak{p}^r$ is Koszul S-module. □

It is a known result that if $\mathfrak{c}$ is $\mathfrak{n}$-primary ideal of S with $\mathfrak{n}^{2r-2} \subseteq \mathfrak{c} \subseteq \mathfrak{n}^r$, then $\mathfrak{c}$ is Golod. This was first noticed by Löfwall [13] also see [7, Example 2.10]. The following extends this result to an ideal generated by a part of a regular system of parameter.

Proposition 2.12. *Let $\mathfrak{p}$ be an ideal of S generated by a part of a regular system of parameter. Assume that r is a positive integer. Then the following holds.*

(1) $\rho(\mathfrak{p}) = 1$;
(2) if $r \geq 2$, then any ideal of S between $\mathfrak{p}^{2r-2}$ and $\mathfrak{p}^r$ is Golod;
(3) if $\mathfrak{a}$ be an ideal of S satisfying $\mathfrak{p}^r \subseteq \mathfrak{a}$, then $\mathfrak{p}^r\mathfrak{a}$ is Golod.

Proof. From Lemma 2.1, $\mathfrak{p}^r$ is Koszul. Hence by Remark 2.9, the map

$$\mathrm{Tor}_i^S(\mathfrak{p}^r, S/\mathfrak{n}^2) \to \mathrm{Tor}_i^S(\mathfrak{p}^r, S/\mathfrak{n})$$

is zero for all $i > 0$. Now, applying similar argument used in the proof of 2.10, one can see that the map

$$\mathrm{Tor}_i^S(\mathfrak{p}^r, S/\mathfrak{n}) \to \mathrm{Tor}_i^S(\mathfrak{p}^{r-1}, S/\mathfrak{n})$$

is zero for all $i > 0$. Thus (1) follows from Remark 2.1. Part (2) follows from (1) and Theorem 2.2. Since $\mathfrak{p}^r$ is Koszul, Theorem 2.10 concludes (3).

□

References

[1] D. J. Anick, A counterexample to a conjecture of Serre, Ann. of Math. 115 (1982), 1-33.

[2] L. L. Avramov, A. R Kustin, M. Miller, Poincaré series of modules over local rings of small embedding codepth or small linking number, J. Algebra 118 (1988), 162-204.

[3] J. Backelin and R. Fröberg, Koszul algebras, Veronese subrings, and rings with linear resolutions, Rev. Roumaine Math. Pures. Appl. 30 (1985), 85-97.

[4] D. Eisenbud, Commutative Algebra with a View Toward Algebraic Geometry, Grad. Text in Math. 150, Springer, New York, 1995.

[5] T. H. Gulliksen, Massey operations and the Poincare series of certain local rings, J. Algebra 22 (1972), 223-232.

[6] J. Herzog, Canonical Koszul cycles, Aportaciones Mat. Notas de Investigacion 6 (1992) 33-41.

[7] J. Herzog, C. Huneke, Ordinary and symbolic powers are Golod, Adv. Math., 246: 89-99 (2013)

[8] J. Herzog, M. Steurich, Golodideale der Gestalt $\mathfrak{a} \cap \mathfrak{b}$, J. Algebra 58 (1979), 31-36.

[9] J. Herzog, S. Iyengar, Koszul modules, J. Pure Appl. Algebra 201 (2005) 154–188.

[10] J. Herzog, V. Welker and S. Yassemi, Homology of powers of ideals: Artin-Rees numbers of syzygies and the Golod property. Preprint (2011), http://arxiv.org/abs/1108.5862.

[11] S. Iyengar, T. Römer, Linearity defects of modules over commutative rings. J. Algebra 322 (2009), 3212-3237.

[12] H. D. Nguyen, Good and bad behaviour of the linearity defect, arXiv:1411.0261

[13] C. Löfwall, On the subalgebra generated by the one-dimensional elements in the Yoneda Ext-algebra, in: Algebra, Algebraic Topology and Their Interactions, (Stockholm, 1983), in: Lecture Notes in Math., vol. 1183, Springer- Verlag, 1983, pp. 291-338.

[14] J.E. Roos, A Gorenstein numerical semi-group ring having a transcendental series of Betti numbers, Preprint, 2012

[15] M.E. Rossi, L.M. Sega, *Poincar series of modules over compressed Gorenstein local rings*, J.Algebra 259 (2014), 421-447.

[16] T. H. Gulliksen, G. Levin, Homology of local rings, Queen's Papers Pure Appl. Math. 20, Queen's Univ., Kingston, ON (1969).

[17] T. Römer, On minimal graded free resolutions, Dissertation, Essen, 2001.

[18] Liana M. Şega, On the linearity defect of the residue field. J. Algebra 384 (2013), 276-290.

[19] J.P. Serre, Algábre Locale et Multiplicits. Lecture Notes in Mathematics, Vol. 11Springer, Berlin (1965).

[20] A. De Stefani, Products of ideals may not be Golod, arXiv:1506.09129.

Rasoul Ahangari Maleki, School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran, Iran

E-mail address: rahangari@ipm.ir, rasoulahangari@gmail.com